



# Predictors of Progression From the Appearance of Islet Autoantibodies to Early Childhood Diabetes: The Environmental Determinants of Diabetes in the Young (TEDDY)

DOI: 10.2337/dc14-2426

Andrea K. Steck,<sup>1</sup> Kendra Vehik,<sup>2</sup> Ezio Bonifacio,<sup>3</sup> Ake Lernmark,<sup>4</sup> Anette-G Ziegler,<sup>5</sup> William A. Hagopian,<sup>6</sup> JinXiong She,<sup>7</sup> Olli Simell,<sup>8</sup> Beena Akolkar,<sup>9</sup> Jeffrey Krischer,<sup>2</sup> Desmond Schatz,<sup>10</sup> and Marian J. Rewers,<sup>1</sup> and the TEDDY Study Group\*

## OBJECTIVE

While it is known that there is progression to diabetes in <10 years in 70% of children with two or more islet autoantibodies, predictors of the progression to diabetes are only partially defined.

## RESEARCH DESIGN AND METHODS

The Environmental Determinants of Diabetes in the Young (TEDDY) study has observed 8,503 children who were at increased genetic risk for autoimmune diabetes. Insulin autoantibodies (IAAs), GAD65 autoantibodies (GADAs), and insulinoma-associated protein 2 autoantibodies (IA-2As) were measured every 3 months until 4 years of age and every 6 months thereafter; if results were positive, the autoantibodies were measured every 3 months.

## RESULTS

Life table analysis revealed that the cumulative incidence of diabetes by 5 years since the appearance of the first autoantibody differed significantly by the number of positive autoantibodies (47%, 36%, and 11%, respectively, in those with three autoantibodies, two autoantibodies, and one autoantibody,  $P < 0.001$ ). In time-varying survival models adjusted for first-degree relative status, number of autoantibodies, age at first persistent confirmed autoantibodies, and HLA genotypes, higher mean IAA and IA-2A levels were associated with an increased risk of type 1 diabetes in children who were persistently autoantibody positive (IAAs: hazard ratio [HR] 8.1 [95% CI 4.6–14.2]; IA-2A: HR 7.4 [95% CI 4.3–12.6];  $P < 0.0001$ ). The mean GADA level did not significantly affect the risk of diabetes.

## CONCLUSIONS

In the TEDDY study, children who have progressed to diabetes usually expressed two or more autoantibodies. Higher IAA and IA-2A levels, but not GADA levels, increased the risk of diabetes in those children who were persistently autoantibody positive.

Autoimmune type 1 diabetes is preceded by a preclinical period characterized by the appearance and persistence of islet insulin autoantibodies (IAAs) (1), GAD autoantibodies (GADAs) (2), insulinoma-associated protein 2 autoantibodies (IA-2As) (3), and zinc transporter 8 autoantibodies (4). Guidelines for screening first-degree

<sup>1</sup>Barbara Davis Center for Childhood Diabetes, University of Colorado School of Medicine, Aurora, CO

<sup>2</sup>Pediatrics Epidemiology Center, University of South Florida, Tampa, FL

<sup>3</sup>Technische Universität Dresden, Dresden, Germany

<sup>4</sup>Lund University/Clinical Research Centre, Skåne University Hospital, Malmö, Sweden

<sup>5</sup>Institute of Diabetes Research, Helmholtz Zentrum München, and Klinikum rechts der Isar, Technische Universität München, and Forschergruppe Diabetes e.V., Neuherberg, Germany

<sup>6</sup>Pacific Northwest Diabetes Research Institute, Seattle, WA

<sup>7</sup>Medical College of Georgia, Georgia Regents University, Augusta, GA

<sup>8</sup>Department of Pediatrics, University of Turku, Turku, Finland

<sup>9</sup>Division of Diabetes, Endocrinology & Metabolic Diseases, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD

<sup>10</sup>University of Florida, Gainesville, FL

Corresponding author: Andrea K. Steck, [andrea.steck@ucdenver.edu](mailto:andrea.steck@ucdenver.edu).

Received 13 October 2014 and accepted 5 January 2015.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/doi:10.2337/dc14-0850/-/DC1>.

\*A complete list of the members of the TEDDY Study Group can be found in the Supplementary Data online.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

relatives (FDRs) of persons with type 1 diabetes exist (5,6); however, nearly 90% of patients in whom type 1 diabetes has been newly diagnosed have no FDRs with the disease. Previous studies (7–10) have estimated that diabetes will develop within 10 years in 27–40% of the general population children expressing two or more autoantibodies. In Finnish children, double positivity for GADA and IA-2A on a one-time screening identified up to 60% of diabetic cases in the ensuing 27 years (11). A recent study (12) combining three prospective cohorts of multiple autoantibody-positive children from Colorado, Finland, and Germany reported a risk of diabetes of 70% within 10 years and 84% within 15 years after seroconversion.

Once persistent islet autoimmunity develops, progression toward diabetes seems to be locked in, although the time to diabetes diagnosis varies tremendously. The determinants of the progression are only partially understood, and include younger age at seroconversion, the number of autoantibodies, and higher levels of IAAs (13).

The Environmental Determinants of Diabetes in the Young (TEDDY) (14) is a multicenter observational study designed to map the events leading to type 1 diabetes from birth to the age of 15 years and to identify the precipitating exposures. In this largest prospective birth cohort of genetically high-risk children, we report predictors of progression from islet autoantibodies to clinical diabetes.

## RESEARCH DESIGN AND METHODS

### Study Population

Since September 2004, TEDDY has accrued and observed a cohort of 8,503 infants who were at increased genetic risk for type 1 diabetes. The vast majority (89%) have no FDR with type 1 diabetes, while 11% are siblings or offspring of a person with type 1 diabetes. The participants were identified at birth through genetic screening for diabetes susceptibility HLA-DR/DQ genotypes at sites in Sweden, Finland, Germany, Colorado, Washington state, and Florida/Georgia. Those participants enrolled in the study are followed up prospectively from birth to 15 years of age, with study visits beginning at 3 months of age, then every 3 months until 4 years of age and every 6 months thereafter. Children who are positive for islet autoantibodies are

followed up every 3 months. The details of screening and follow-up have been previously published (15,16). A total of 577 children in whom persistent confirmed islet autoimmunity developed were included in this study; 164 of those children progressed to diabetes. We did not include children who were not positive for antibodies prior to diagnosis ( $N = 12$ ). Subjects with positive but not persistent islet autoantibodies ( $N = 505$ ), subjects who had persistent confirmed autoantibodies but who withdrew from the study ( $N = 28$ ), or had maternal autoantibodies ( $N = 198$ ) were excluded from these analyses. The study was approved at all sites by local institutional review boards.

### Islet Autoantibodies

GADAs, IA-2As, and IAAs were measured in two laboratories by standard radio-binding assays (17–19). For sites in the U.S., all serum samples were assayed at the Barbara Davis Center for Childhood Diabetes at the University of Colorado at Denver. In Europe, all sera were assayed at the University of Bristol (Bristol, U.K.). Both laboratories have previously shown high assay sensitivity and specificity as well as concordance (20). All results that were positive for islet autoantibodies and 5% of negative samples were retested in the other reference laboratory and deemed confirmed if results were concordant. In addition, the TEDDY study participated in the National Institute of Diabetes and Digestive and Kidney Diseases harmonization project, in collaboration with the Diabetes Research Institute (Munich, Germany), to evaluate the impact of the harmonized assay protocol on the concordance of IA-2A and GADA results (17,21). On the retested TEDDY study samples, discordance decreased from 4% to 1.8% for IA-2As ( $n = 604$  samples) and from 15.4% to 2.7% for GADAs ( $n = 515$  samples). Persistent islet autoantibodies were defined as being confirmed positive GADA, IA-2A, or IAA on at least two consecutive study visits. Transplacental autoantibodies (defined as the transient presence of the same autoantibody in a child younger than 18 months of age and his/her mother) were excluded from analyses as were positive or confirmed but not persistent confirmed autoantibodies. Fluctuating positive autoantibodies were defined as being persistently confirmed positive

autoantibodies in subjects whose samples reverted to negativity at one or more time points. Nonfluctuating persistent islet autoantibodies were defined as confirmed autoantibody positivity for a specific autoantibody on two consecutive visits and did not revert to negative after initial detection (i.e., stayed positive throughout follow-up). Diabetes was defined according to American Diabetes Association criteria for diagnosis (22).

### Statistical Analysis

Data were analyzed using the Statistical Analysis System Software (version 9.2; SAS Institute, Cary, NC). Categorical variables were analyzed using Pearson  $\chi^2$  tests. Continuous variables were tested using the  $t$  test for differences in means or the Wilcoxon rank sum test for differences in medians. Due to different autoantibody cutoff values in Bristol and Denver, and change in the type of assay over time from the TEDDY assay to the harmonized assay (20% of the enrolled population is missing a harmonized measure prior to January 2010), autoantibody levels were converted to SD units away from threshold (“z scores”) for analysis. For IA-2As and GADAs, the harmonized assay was used; if a measure was missing from the harmonized assay, the TEDDY assay was used (the Bristol z score was used and, if missing, then the Denver z score was used). As for IAAs (not harmonized), the Bristol z score was used, and, if missing, the Denver z score was used. The IAA, GADA, and IA-2A levels were log transformed for analyses. Kaplan-Meier life tables were used to determine the time to diabetes onset from the initial seroconversion for each subject and were compared using the log-rank  $\chi^2$  statistic.

Multivariate time-varying Cox proportional hazards models were used to estimate the risk of diabetes and to determine significant predictors; country of residence was the stratified variable. Three time-varying covariates were considered (mean autoantibody level for IA-2As, IAAs, and GADAs). Risk periods were taken as the time between serum sample collections at consecutive (3-month) visits where subjects were still deemed to be at risk for diabetes from the time of persistent confirmed autoantibody positivity to the diagnosis of diabetes or last serum sample collection. Time to diabetes was defined as

the time from initial seroconversion. The Efron method for tied survival times was used in the Cox analysis and the Akaike information criterion to assess the best model. The counting process format in SAS version 9.3 was applied to fit the time-dependent Cox proportional hazard models. HLA effect was assessed in the time-varying Cox models as the highest-risk HLA-DR3-DQ2/DR4-DQ8 genotype versus other genotypes as well as by evaluating the following five primary TEDDY eligibility genotypes groups: HLA-DR3/4, DR4/4, DR3/3, DR4/8, and additional eligibility genotypes applied only to FDRs (HLA-DR3/9, DR4/1, DR4/4b, DR4/9, and DR4/13) (16). The non-time-varying variables (age at first persistent confirmed autoimmunity, FDR status, number of autoantibodies when first positive, and HLA genotype) were first evaluated as predictors of progression to type 1 diabetes, then the time-varying variables (mean log micro-IAA [mIAA], mean log GADA, and mean log IA-2A) were added.

In the subpopulation in whom diabetes developed, general linear models were used to evaluate potential predictors of age of diabetes onset; stepwise selection using the Cp statistic was used to assess best fit. A two-tailed *P* value with an  $\alpha$  level for significance was set at 0.05.

## RESULTS

During a median follow-up time of 5.0 years (interquartile range 2.6–6.6

years), persistent confirmed islet autoimmunity developed in 577 children, 164 of whom have progressed to diabetes. In those with persistent islet autoantibodies, 61% had multiple autoantibodies, including 32% with all three autoantibodies present, 17% with IAAs and GADAs, 7% with IAAs and IA-2As, and 5% with GADAs and IA-2As. Of those with a single confirmed persistent antibody, 22% had GADAs, 16% had IAAs, and 1% had IA-2As. The age of seroconversion was significantly younger in subjects who progressed to diabetes compared with those who did not (1.3 vs. 2.5 years, respectively,  $P < 0.0001$ ) and varied by autoantibody type (Table 1). Among subjects who progressed to diabetes ( $n = 164$ ), the great majority (88%) had multiple autoantibodies, including 54% with all three autoantibodies, 16% with IAAs and GADAs, 15% with IAAs and IA-2As, and 3% with GADAs and IA-2As. Of those who progressed to diabetes, 9% had mIAAs alone, 2% had GADAs alone, and 1% had IA-2As alone.

Progression to diabetes by Kaplan-Meier life table analysis (Fig. 1) revealed that the cumulative incidence of diabetes within 5 years of seroconversion increased with the number of positive autoantibodies, 47%, 36%, and 11%, in those with, respectively three autoantibodies, two autoantibodies, and one autoantibody ( $P < 0.0001$ ). There was

no significant difference between children expressing two or three autoantibodies ( $P = 0.08$ ). FDRs had an increased 5-year risk of progression to diabetes, compared with children without a close relative with diabetes, in those with one autoantibody (21% vs. 9%,  $P = 0.04$ ) or two autoantibodies (54% vs. 30%;  $P = 0.01$ ), but not in those with three autoantibodies (54% vs. 45%,  $P = 0.17$ ) (Fig. 2). There was no difference in 5-year progression to diabetes by high-risk HLA-DR3-DQ2/DR4-DQ8 genotype in children expressing one autoantibody (16% vs. 7%,  $P = 0.08$ ), two autoantibodies (37% vs. 35%,  $P = 0.85$ ), or three autoantibodies (48% vs. 47%,  $P = 0.66$ ) (Supplementary Fig. 1).


We compared progression to diabetes in children with persistent and fluctuating islet autoantibodies (Supplementary Fig. 2). Subjects with persistent IAAs had an increased risk of progression to diabetes compared with subjects with fluctuating IAA levels ( $<0.001$ ); the estimated proportion of children progressing to diabetes at 5 years was 57% versus 22%, respectively. In contrast, there was no difference in the progression to diabetes between children with fluctuating versus persistent IA-2As ( $P = 0.42$ ) or GADAs ( $P = 0.53$ ).

In multivariate time-varying Cox proportional hazards models, higher IAA and IA-2A levels were significant predictors of type 1 diabetes in children who

**Table 1—Characteristics of TEDDY subjects by antibody and diabetes status**

Characteristic	Persistent confirmed Ab+ (N = 577)	Ab– (N = 7,195)	<i>P</i> value (Ab+ vs. Ab–)	Persistent confirmed Ab+ (no diabetes) (N = 413)	Type 1 diabetes* (N = 164)	<i>P</i> value (Ab+ vs. diabetes)†
Male sex, <i>n</i> (%)	326 (57)	3,621 (50)	0.0043	236 (57)	90 (55)	0.6206
FDR (%)	124 (21)	586 (8)	<0.0001	73 (18)	51 (31)	0.0004
HLA-DR3/4	293 (51)	2,747 (38)	<0.0001	200 (48)	93 (57)	0.0446
HLA-DR4/4	100 (17)	1,406 (20)		77 (19)	23 (14)	
HLA-DR3/3	72 (12)	1,561 (22)		57 (14)	15 (9)	
HLA-DR4/8	86 (15)	1,294 (18)		65 (16)	21 (13)	
Other HLA genotypes‡	26 (5)	187 (3)		14 (3)	12 (7)	
Follow-up duration (years)	5.73 (4.4–7.2)	4.81 (2.1–6.6)	<0.0001	6.36 (5.1–7.5)	3.35 (2.1–5.2)	<0.0001
Age at first Ab positivity	2.06 (1.2–3.4)			2.52 (1.5–4.0)	1.28 (0.9–2.0)	<0.0001
First mIAAs (years)	1.96 (1.1–3.2)			2.46 (1.3–3.6)	1.29 (1.0–2.0)	<0.0001
First GADs (years)	2.48 (1.5–3.8)			3.02 (2.0–4.3)	1.54 (1.1–2.3)	<0.0001
First IA-2As (years)	2.56 (1.8–4.0)			3.58 (2.5–4.9)	2.00 (1.3–2.7)	<0.0001
Age at diabetes diagnosis (years)					3.30 (2.0–5.1)	

Data are reported as *n* (%) or median (quartile 1–quartile 3), unless otherwise indicated. Ab–, antibody negative; Ab+, antibody positive. \*Diabetic subjects are included within the Persistent confirmed Ab+ ( $n = 577$ ) population. †Comparison between those Ab+ subjects who did not progress to type 1 diabetes and those in whom diabetes developed. ‡FDR-specific HLA genotypes.



**Figure 1**—Progression to diabetes in children with confirmed autoantibodies ( $N = 577$ ). Ab+, antibody positive.

were persistently autoantibody positive for one or more islet autoantibodies. A 1-unit increase in log mean IAA level increased the diabetes risk approximately eightfold (hazard ratio [HR] 8.12 [95% CI 4.6–14.2],  $P < 0.0001$ ), and a 1-unit increase in log mean IA-2A level increased diabetes risk  $\sim 7.5$ -fold (HR 7.4 [95% CI 4.3–12.6],  $P < 0.0001$ ) (Table 2) after adjustment of FDR status

(HR 1.6 [95% CI 1.1–2.4],  $P = 0.02$ ), age at first persistent confirmed autoantibodies (HR 0.97 [95% CI 0.92–0.98],  $P < 0.0001$ ), and the number of autoantibodies when first confirmed positive (two antibodies vs. one antibody: HR 1.5 [95% CI 1.0–2.4],  $P = 0.046$ ; three antibodies vs. one antibody: HR 1.7 [95% CI 0.8–3.7],  $P = 0.21$ ). HLA genotype ( $P = 0.11$ ) and log mean GADA level


( $P = 0.85$ ) did not significantly affect the risk of diabetes.

In subjects in whom diabetes developed ( $N = 164$ ), general linear models were used to explore predictors of the age of diabetes onset. The number of initial autoantibodies, mean IAA and IA-2A levels, initial IA-2A level, country of residence, and age of first seroconversion provided the best predictive model in estimating the age of diabetes onset ( $r = 0.56$ ,  $P < 0.0001$ ).

### CONCLUSIONS

This large prospective cohort study addressed the critical challenge of predicting the absolute risk and the time to progression to clinical diabetes among children in whom persistent islet autoimmunity developed. The majority of children who progressed to diabetes had two or more islet autoantibodies and were very young at seroconversion (median 1.3 years). In children with confirmed positive autoantibodies, progression to diabetes was influenced by family history, but not by the presence of the high-risk HLA-DR3-DQ2/DR4-DQ8 genotype. The high risk of progression to diabetes in children with multiple islet autoantibodies has previously been shown in individual studies (13,23,24) as well as in a recent study (12) combining three prospective cohorts from Colorado, Finland, and Germany. In DAISY, the high-risk HLA-DR3-DQ2/DR4-DQ8 genotype influenced progression to type 1 diabetes in children expressing two or fewer positive islet autoantibodies, but not in those expressing three positive islet autoantibodies (13). On the other hand, family history did not influence progression to diabetes in children with two or more autoantibodies. These discrepancies between previous reports and TEDDY results are most likely due to differences in the duration of follow-up, although differences in autoantibody assays or slight differences in HLA inclusion criteria cannot be excluded.

The TEDDY study is currently the largest prospective study with very detailed genetic and antibody follow-up data from birth of infants with high-risk genes for type 1 diabetes, including both general population children and FDRs. The frequent antibody testing in this young cohort may improve the accuracy of the time to event analyses.



**Figure 2**—Progression to diabetes in children expressing 1 autoantibody, 2 autoantibodies or 3 autoantibodies by family history. Ab+, antibody positive; GP, general population.

**Table 2—Multivariate time-varying Cox proportional hazards models**

	Adjusted HR	95% CI	P value
Mean mIAAs	8.12	4.64–14.23	<0.0001
Mean IA-2As	7.40	4.34–12.55	<0.0001
Mean GADAs	1.06	0.62–1.80	0.845
FDR	1.58	1.07–2.35	0.023
Age at first persistent confirmed Ab+	0.97	0.92–0.98	<0.0001
Number of Abs at first confirmed Ab+			
2 Abs vs. 1 Ab	1.54	1.01–2.35	0.046
3 Abs vs. 1 Ab	1.67	0.75–3.71	0.212
HLA genotype			
DR3/4 vs. DR4/4	1.37	0.82–2.29	0.227
DR3/3 vs. DR4/4	2.71	1.29–5.68	0.008
DR4/8 vs. DR4/4	1.14	0.61–2.13	0.689
FDR-specific† vs. DR4/4	1.36	0.60–3.08	0.457

Ab, antibody. †FDR specific HLA: HLA DR3/9, DR4/1, DR4/4b, DR4/9 and DR4/13.

Progression toward diabetes seems to be locked in once persistent islet autoimmunity develops; however, the time to diabetes diagnosis varies tremendously, and the factors influencing progression to diabetes are still poorly understood. In this TEDDY cohort of subjects with persistent confirmed islet autoimmunity, higher mean IAA and IA-2A levels were associated with an increased risk of type 1 diabetes in time-varying survival models adjusted for FDR status, number of autoantibodies, age at first persistent confirmed autoantibodies, and HLA genotypes. In addition, in subjects in whom diabetes developed, predictors of the age of diabetes onset included the number of initial autoantibodies, mean IAA and IA-2A levels, initial IA-2A level, country of residence, and age at first seroconversion, confirming that it is possible to estimate the age of diabetes onset, as shown previously in a smaller prospective study (13).

Despite a limited follow-up duration of only 5 years, our results support the notion that diabetes is likely to develop in most children with persistent multiple islet autoantibodies. However, the time of progression to diabetes can be highly variable. Therefore, factors that can predict the age at development of diabetes may have an important prognostic value for parents and providers, and may also be of use in interim analyses of prevention trials. In the first 100 children in the TEDDY study in whom diabetes developed, FDRs had a higher cumulative incidence than children from the general population (25), but the FDR status did not affect the progression to diabetes in the present analysis once

three or more autoantibodies had developed. While the cumulative risk of diabetes can vary according to age, relationship to the proband, positivity for IAA, IA-2A, and GADA, the number and combination of islet autoantibodies, HLA-DR/DQ genotype, baseline glucose tolerance, and first-phase insulin secretion (26–31), the TEDDY study results narrow down the list of predictors to young age at seroconversion, positivity for multiple autoantibodies, high autoantibody levels, and persistent positivity for IAA. In this regard, our results are consistent with those of prior studies (13,32,33) but include more participants and more frequent antibody testing, which may improve the accuracy of the time-to-event analyses. Only IAA and IA-2A levels (and not GADA levels) were useful in estimating the age at diabetes onset. Both nonfluctuating IAA and IA-2A levels were significant factors for progression to diabetes, confirming findings from a previous smaller prospective study (13). The mechanism underlying the specific association of levels of IAAs with the rate of progression to diabetes is not defined but may relate to the hypothesized unique biologic importance of insulin autoimmunity  $\beta$ -cell destruction. On the other hand, the IAA is most often the initial antibody to become positive in young children; this TEDDY cohort is still very young, and these antibody findings might be different in an older population.

TrialNet and other studies (34,35) have demonstrated a period of impaired fasting glucose levels or impaired glucose tolerance that precedes type 1 diabetes onset by several months or years

among persons positive for islet autoantibodies. The oral glucose tolerance test performed in clinical trials of type 1 diabetes prevention has long been known to have value in predicting progression to diabetes among subjects with islet autoantibodies (34). The DPT-1 (Diabetes Prevention Trial-Type 1) study (36,37) in FDR subjects reported that a risk score based on age, BMI, and oral glucose tolerance test indexes (glucose and C-peptide values), without the use of intravenous glucose tolerance tests or additional autoantibodies, accurately predicted diabetes risk over a short time period in islet cell antibody-positive relatives. The likelihood of progression to diabetes increased with mild fasting or after oral glucose load dysglycemia. However, the interval between dysglycemia and clinical diabetes is often very short, suggesting that the best opportunity to prevent type 1 diabetes is before this “dysglycemic” period, and must rely on screening for immunogenetic markers (38) such as HLA and islet autoantibodies, which are extensively measured in the TEDDY study. Further studies will be required to confirm the predictive value of these antibody tests and their combinations for clinical diabetes appearing later in childhood, as well as for children in whom autoantibodies first develop after puberty.

**Funding.** The TEDDY Study Group is funded by grants U01-DK-63829, U01-DK-63861, U01-DK-63821, U01-DK-63865, U01-DK-63863, U01-DK-63836, U01-DK-63790, UC4-DK-63829, UC4-DK-63861, UC4-DK-63821, UC4-DK-63865, UC4-DK-63863, UC4-DK-63836, and UC4-DK-95300, and by Contract no. HHSN267200700014C from the National Institute of Diabetes and Digestive and Kidney Diseases, National Institute of Allergy and Infectious Diseases, National Institute of Child Health and Human Development, National Institute of Environmental Health Sciences, JDRF, and Centers for Disease Control and Prevention. This work was supported in part by the National Institutes of Health/National Center for Advancing Translational Sciences Clinical and Translational Science Awards to the University of Florida (UL1-TR-000064) and the University of Colorado (UL1-TR-001082).

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** A.K.S. wrote the manuscript and contributed to the discussion. K.V. and E.B. researched the data, and reviewed and edited the manuscript. A.L., W.A.H., B.A., J.K., D.S., and M.J.R. contributed to the discussion, and reviewed and edited manuscript. A-G.Z., J.S., and O.S. reviewed and edited the manuscript. A.K.S. is the guarantor of this work and,

as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- Palmer JP, Asplin CM, Clemons P, et al. Insulin antibodies in insulin-dependent diabetics before insulin treatment. *Science* 1983; 222:1337–1339
- Baekkeskov S, Aanstoot H-J, Christgau S, et al. Identification of the 64K autoantigen in insulin-dependent diabetes as the GABA-synthesizing enzyme glutamic acid decarboxylase. *Nature* 1990;347:151–156
- Gianani R, Rabin DU, Verge CF, et al. ICA512 autoantibody radioassay. *Diabetes* 1995;44: 1340–1344
- Wenzlau JM, Juhl K, Yu L, et al. The cation efflux transporter ZnT8 (Slc30A8) is a major autoantigen in human type 1 diabetes. *Proc Natl Acad Sci U S A* 2007;104:17040–17045
- Bingley PJ, Bonifacio E, Ziegler AG, Schatz DA, Atkinson MA, Eisenbarth GS; Immunology of Diabetes Society. Proposed guidelines on screening for risk of type 1 diabetes. *Diabetes Care* 2001;24:398
- Verge CF, Gianani R, Kawasaki E, et al. Prediction of type 1 diabetes in first-degree relatives using a combination of insulin, GAD, and ICA512/bdc/IA-2 autoantibodies. *Diabetes* 1996; 45:926–933
- LaGasse JM, Brantley MS, Leech NJ, et al.; Washington State Diabetes Prediction Study. Successful prospective prediction of type 1 diabetes in schoolchildren through multiple defined autoantibodies: an 8-year follow-up of the Washington State Diabetes Prediction Study. *Diabetes Care* 2002;25:505–511
- Maclaren NK, Lan MS, Schatz D, Malone J, Notkins AL, Krischer J. Multiple autoantibodies as predictors of type 1 diabetes in a general population. *Diabetologia* 2003;46:873–874
- Bingley PJ, Bonifacio E, Williams AJ, Genovese S, Bottazzo GF, Gale EA. Prediction of IDDM in the general population: strategies based on combinations of autoantibody markers. *Diabetes* 1997;46:1701–1710
- Samuelsson U, Sundkvist G, Borg H, Fernlund P, Ludvigsson J. Islet autoantibodies in the prediction of diabetes in school children. *Diabetes Res Clin Pract* 2001;51:51–57
- Knip M, Korhonen S, Kulmala P, et al. Prediction of type 1 diabetes in the general population. *Diabetes Care* 2010;33:1206–1212
- Ziegler AG, Rewers M, Simell O, et al. Seroprevalence to multiple islet autoantibodies and risk of progression to diabetes in children. *JAMA* 2013;309:2473–2479
- Steck AK, Johnson K, Barriga KJ, et al. Age of islet autoantibody appearance and mean levels of insulin, but not GAD or IA-2 autoantibodies, predict age of diagnosis of type 1 diabetes: diabetes autoimmunity study in the young. *Diabetes Care* 2011;34:1397–1399
- TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study. *Ann N Y Acad Sci* 2008;1150:1–13
- Kiviniemi M, Hermann R, Nurmi J, et al.; TEDDY Study Group. A high-throughput population screening system for the estimation of genetic risk for type 1 diabetes: an application for the TEDDY (the Environmental Determinants of Diabetes in the Young) study. *Diabetes Technol Ther* 2007;9:460–472
- Hagopian WA, Erlich H, Lernmark A, et al.; TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY): genetic criteria and international diabetes risk screening of 421 000 infants. *Pediatr Diabetes* 2011;12:733–743
- Bonifacio E, Yu L, Williams AK, et al. Harmonization of glutamic acid decarboxylase and islet antigen-2 autoantibody assays for National Institute of Diabetes and Digestive and Kidney Diseases consortia. *J Clin Endocrinol Metab* 2010;95:3360–3367
- Babaya N, Yu L, Miao D, et al. Comparison of insulin autoantibody: polyethylene glycol and micro-IAA 1-day and 7-day assays. *Diabetes Metab Res Rev* 2009;25:665–670
- Yu L, Robles DT, Abiru N, et al. Early expression of antiinsulin autoantibodies of humans and the NOD mouse: evidence for early determination of subsequent diabetes. *Proc Natl Acad Sci U S A* 2000;97:1701–1706
- Törn C, Mueller PW, Schlosser M, Bonifacio E, Bingley PJ. Diabetes Antibody Standardization Program: evaluation of assays for autoantibodies to glutamic acid decarboxylase and islet antigen-2. *Diabetologia* 2008;51:846–852
- Vehik K, Fiske SW, Logan CA, et al.; TEDDY Study Group. Methods, quality control and specimen management in an international multicentre investigation of type 1 diabetes: TEDDY. *Diabetes Metab Res Rev* 2013;29:557–567
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2014;37(Suppl. 1):S81–S90
- Ziegler AG, Hummel M, Schenker M, Bonifacio E. Autoantibody appearance and risk for development of childhood diabetes in offspring of parents with type 1 diabetes: the 2-year analysis of the German BABYDIAB study. *Diabetes* 1999;48:460–468
- Kukko M, Kimpimäki T, Korhonen S, et al. Dynamics of diabetes-associated autoantibodies in young children with human leukocyte antigen-conferred risk of type 1 diabetes recruited from the general population. *J Clin Endocrinol Metab* 2005;90:2712–2717
- Elding Larsson H, Vehik K, Gesualdo P, et al.; TEDDY Study Group. Children followed in the TEDDY study are diagnosed with type 1 diabetes at an early stage of disease. *Pediatr Diabetes* 2014;15:118–126
- Bingley PJ, Gale EA; European Nicotinamide Diabetes Intervention Trial (ENDIT) Group. Progression to type 1 diabetes in islet cell antibody-positive relatives in the European Nicotinamide Diabetes Intervention Trial: the role of additional immune, genetic and metabolic markers of risk. *Diabetologia* 2006;49:881–890
- Mrena S, Virtanen SM, Laipala P, et al. Models for predicting type 1 diabetes in siblings of affected children. *Diabetes Care* 2006;29: 662–667
- Orban T, Sosenko JM, Cuthbertson D, et al.; Diabetes Prevention Trial-Type 1 Study Group. Pancreatic islet autoantibodies as predictors of type 1 diabetes in the Diabetes Prevention Trial-Type 1. *Diabetes Care* 2009;32:2269–2274
- Walter M, Albert E, Conrad M, et al. IDDM2/insulin VNTR modifies risk conferred by IDDM1/HLA for development of type 1 diabetes and associated autoimmunity. *Diabetologia* 2003; 46:712–720
- Steck AK, Bugawan TL, Valdes AM, et al. Association of non-HLA genes with type 1 diabetes autoimmunity. *Diabetes* 2005;54:2482–2486
- Bonifacio E, Hummel M, Walter M, Schmid S, Ziegler AG. IDDM1 and multiple family history of type 1 diabetes combine to identify neonates at high risk for type 1 diabetes. *Diabetes Care* 2004;27:2695–2700
- Siljander HT, Simell S, Hekkala A, et al. Predictive characteristics of diabetes-associated autoantibodies among children with HLA-conferred disease susceptibility in the general population. *Diabetes* 2009;58:2835–2842
- Achenbach P, Warncke K, Reiter J, et al. Stratification of type 1 diabetes risk on the basis of islet autoantibody characteristics. *Diabetes* 2004;53:384–392
- Sosenko JM, Palmer JP, Greenbaum CJ, et al.; Diabetes Prevention Trial-Type 1 Study Group. Increasing the accuracy of oral glucose tolerance testing and extending its application to individuals with normal glucose tolerance for the prediction of type 1 diabetes: the Diabetes Prevention Trial-Type 1. *Diabetes Care* 2007;30:38–42
- Stene LC, Barriga K, Hoffman M, et al. Normal but increasing hemoglobin A1c levels predict progression from islet autoimmunity to overt type 1 diabetes: Diabetes Autoimmunity Study in the Young (DAISY). *Pediatr Diabetes* 2006;7:247–253
- Sosenko JM, Krischer JP, Palmer JP, et al.; Diabetes Prevention Trial-Type 1 Study Group. A risk score for type 1 diabetes derived from autoantibody-positive participants in the diabetes prevention trial-type 1. *Diabetes Care* 2008;31:528–533
- Sosenko JM, Skyler JS, Mahon J, et al.; Type 1 Diabetes TrialNet Study Group; Diabetes Prevention Trial-Type 1 Study Group. The application of the diabetes prevention trial-type 1 risk score for identifying a preclinical state of type 1 diabetes. *Diabetes Care* 2012;35:1552–1555
- Schlosser M, Strebelow M, Wassmuth R, et al. The Karlsburg type 1 diabetes risk study of a normal schoolchild population: association of beta-cell autoantibodies and human leukocyte antigen-DQB1 alleles in antibody-positive individuals. *J Clin Endocrinol Metab* 2002;87: 2254–2261